

RU M 7500.000.002

Visceralgine™

Injection solution in ampoules

Composition

Each 2 ml aqueous solution contains:
Tiemonium Methylsulphate INN 5 mg and
Sodium Chloride BP 18 mg

Therapeutic Indications

This drug is an antispasmodic : It reduces muscle spasms of the intestine, biliary system, bladder and uterus.

This drug is indicated for acute pain.

Contra-Indications

This drug SHOULD NOT BE USED in the following cases :

- glaucoma (acute pain of eyeball with vision disturbances)
- difficulty to urinate (disorders of prostate or bladder)

IN CASE OF DOUBT, YOU MUST ASK YOUR PHYSICIAN FOR ADVICE.

Dosage and Method of Administration

In adults the usual dosage is 1 ampoule by slow IM or IV injection 3 times/24h.

IN ALL CASES, FOLLOW YOUR PHYSICIAN'S PRESCRIPTION TO THE LETTER.

Precautions for use

IN ORDER TO AVOID POSSIBLE DRUG INTERACTIONS, YOU MUST SYSTEMATICALLY INFORM YOUR PHYSICIAN ABOUT ANY OTHER DRUGS YOU ARE TAKING.

Avoid taking this drug without medical advice during pregnancy or breast feeding.

IN CASE OF DOUBT, DO NOT HESITATE TO SEEK ADVICE FROM YOUR PHYSICIAN. KEEP OUT OF THE REACH OF CHILDREN.

Other Possible Effects of the drug

LIKE ANY ACTIVE DRUG, THIS MEDICATION MAY HAVE SOME UNDESIRABLE EFFECTS IN CERTAIN INDIVIDUALS.

Risk of hypotension and tachycardia in the case of rapid injection.

Storage Condition

Store in a dry & dark place in between 2°-30°C. Keep out of the reach of children.

Commercial Pack

Each box contains 10 x 2 ml ampoules in PVC tray.



Manufactured by:
Nuvista Pharma Limited
Tongi - Gazipur

3056DS1401